

2023

FACIES AND DEPOSITIONAL ENVIRONMENTS OF THE MIOCENE SEDIMENTARY SEQUENCE IN NORTH OF TORBALI, IZMIR

Uğur İNCİ
mta@mta.gov.tr

Follow this and additional works at: <https://bmta.researchcommons.org/journal>

 Part of the [Geology Commons](#)

Recommended Citation

İNCİ, Uğur (2023) "FACIES AND DEPOSITIONAL ENVIRONMENTS OF THE MIOCENE SEDIMENTARY SEQUENCE IN NORTH OF TORBALI, IZMIR," *Bulletin of the Mineral Research and Exploration*: Vol. 1991: Iss. 112, Article 2.

Available at: <https://bmta.researchcommons.org/journal/vol1991/iss112/2>

This Research Articles is brought to you for free and open access by Bulletin of the Mineral Research and Exploration. It has been accepted for inclusion in Bulletin of the Mineral Research and Exploration by an authorized editor of Bulletin of the Mineral Research and Exploration.

FACIES AND DEPOSITIONAL ENVIRONMENTS OF THE MIOCENE SEDIMENTARY SEQUENCE IN NORTH OF TORBALI, IZMİR

Ugur İNCİ**

ABSTRACT— A Late Miocene sedimentary sequence, over 900 m., crops out in the north-northeast trending graben type depression developed at the western margin (north of Torbalı) of the Bozdağ high. This sequence is represented with the several complete and incomplete sedimentary cycles which are mainly composed of, in ascending order, conglomerate, sandstone and algal limestone segments. According to the lithofacies analyses, the clastic rocks of this normal fault-bounded depression deposited in the alluvial fan and plain environmental conditions by rapid deposition of the sedimentary load of the gravel-sand dominated hyperconcentrated flood flows originated from the Nif Dağı and Bozdağ high. In consequence of these floods, the small, variable bounded, occasionally wavered and fringing with swamps, ephemeral freshwater lake environments formed characterizing with algal limestones.