

2023

## POSSIBLE PAN-AFRICAN METAVOLCANICS IN THE ÖDEMiŞ, SUBMASSIF OF THE MENDERES MASSIF, WESTERN TURKEY

Osman CANDAN  
osman.candan@deu.edu.tr

Nejat KUN  
nejat.kun@deu.edu.tr

Follow this and additional works at: <https://bmta.researchcommons.org/journal>



Part of the [Earth Sciences Commons](#)

---

### Recommended Citation

CANDAN, Osman and KUN, Nejat (2023) "POSSIBLE PAN-AFRICAN METAVOLCANICS IN THE ÖDEMiŞ, SUBMASSIF OF THE MENDERES MASSIF, WESTERN TURKEY," *Bulletin of the Mineral Research and Exploration*: Vol. 1991: Iss. 112, Article 3.

Available at: <https://bmta.researchcommons.org/journal/vol1991/iss112/3>

This Research Articles is brought to you for free and open access by Bulletin of the Mineral Research and Exploration. It has been accepted for inclusion in Bulletin of the Mineral Research and Exploration by an authorized editor of Bulletin of the Mineral Research and Exploration.

## **POSSIBLE PAN-AFRICAN METAVOLCANICS IN THE ÖDEMiŞ, SUBMASSIF OF THE MENDERES MASSIF, WESTERN TURKEY**

Osman CANDAN\* and Nejat KUN\*

**ABSTRACT.** - Menderes massif located in the Western Anatolia, in Turkey, is made up of the old crystalline rocks. The last main metamorphism of this massif has been completed during Upper Paleocene - Lower Eocene. The general sequence of the study area which is situated in the Ödemiş submassif of the Menderes massif consists of, in ascending order, basal gneiss complex, blue augen gneiss, leptite (metavolcanic), schist and marble. This metamorphic sequence was cut by the post-metamorphic acidic and basic plutons and covered by the young volcanics. Leptites occurring between gneiss and schist units are the metamorphic equivalents of the dacitic and rhyolitic volcanics. These island arc volcanics have a calc-alkaline kindred. According to the age, geological position and chemical characteristics, these metavolcanics of the Ödemiş submassif may be correlated with the island arc volcanic belt widely exposed in the NE Africa and Arabian Peninsula, in relation to the late phases of the Pan-African orogenesis.

### **INTRODUCTION**

Menderes massif consists of the old crystalline rocks which are exposed widely in Western Anatolia, in Turkey. The previous studies began with Phillipson (1911) and continued later by various investigators until today. The presence of some different rock types from the others such as gneiss, schist and phyllite in the Menderes massif was first noted by Schuiling (1962). It was determined by Kun (1983) that these rocks exposed in the Çine submassif of the Menderes massif were clearly originated from the old volcanic rocks and they were called as leptite by the same author based on the similar chemical properties, with those of in Sweden.

In recent years, the relation between the volcanics of the Pan-African orogenesis exposed in NE Africa and Arabian Peninsula and the metavolcanics which show widespread distribution in the Menderes massif has been the subject of investigation by many investigators. In this paper, an attempt is made to solve the problem of the origin and geotectonic position of the leptites in the Ödemiş submassif of the Menderes massif in Turkey.

### **LITHOSTRATIGRAPHY**

The study area is located in the Ödemiş submassif of the Menderes massif, in Western Anatolia in Turkey (Fig. 1 A-B). The generalized columnar section of this area can be given as follows: The lowest level of the rock succession consist of a basal gneiss complex made up of migmatite, augen, granitic and banded gneisses (Fig. 2). In these rocks which contain almandine-rich garnet, the feldspar porphyroblasts amount up to 7-8 cm in length. Through the microscobic studies, the general mineral assemblage of the gneisses which show widely cataclastic texture, was found as "Quartz-plagioclase (oligoclase)-K. feldspar (orthoclase and microcline)-biotite-muscovite-garnet-tourmaline-zircon and apatite".

The blue augen gneisses which have been firstly preserved in the Menderes massif, are found near the contacts of the leptites and basal gneisses as interlayers and lenses in both unit with different lengths. This kind of gneisses typified by the very large blue-violet K-feldspar porphyroblasts, show great similarity to the basal gneiss complex occurring at the bottom of the sequence in terms of mineralogical composition. But, by contrast with basal gneisses, sillimanite is accompanied to garnet in these rocks. The blue-augen gneisses containing abundant relicts of leptite are apparently related to the volcanic origin.

The leptite which lies conformably on the gneisses, vary in thickness from 20-25 m to several kilometers. Some key minerals, such as sillimanite and kyanite, are found in these metavolcanics. The leptites are overlain by the schists with an

\* Dokuz Eylül Üniversitesi, Jeoloji Mühendisliği Bölümü, İzmir-Turkey.




Fig. 1 A - Location map of the study area.

apparently conformable contact. The schist series starts with the kyanite-staurolite-garnet schist at the base and passes to the garnet-mica schists towards the upper level.

Phyllite, marble and muscovite-quartz schists occur as interlayers and lenses in this schist sequence with highly variable thicknesses. Coarse-grained kyanite-staurolite-garnet schists contain widely distributed kyanite-pegmatoid lenses which are parallel to the schistosity. This schist unit consist of "Quartz-plagioclase-kyanite-staurolite-garnet-biotite-muscovite-chlorite-rutil-apatite and zircon". Also, garnet-mica schists are mainly composed of "Quartz-plagioclase-biotite-muscovite-chlorite-garnet-apatite-tourmaline and zircon". The phyllites which are observed as interlayers and lenses show a number of different mineral paragenesis as follows:

Quartz-muscovite-garnet-chloritoid;

Quartz-muscovite-garnet-chloritoid-staurolite;

Quartz-muscovite-garnet-staurolite;

Quartz-muscovite-garnet-staurolite-kyanite.




Fig. 1 B - Geological map of the study area.

The marble interlayers become dominant at the upper level of the metamorphic sequence and these rocks form thick calcareous series at the NE part of the study area.




Fig. 2 - Generalized columnar section of the Ödemiş submassif.


All these metamorphic units were cut by the unmetamorphosed acidic and basic plutons and covered by the Neogene aged volcanic rocks. The acidic magmatites which are of granitic and granodioritic composition, are exposed in gneisses and leplites as stocks with different sizes. The general mineral composition of these rocks were determined as

"Quartz+plagioclase+orthoclase+biotite+muscovite+apatite+zircon±sillimanite and ±gamet". The basic plutons, olivine gabbros, were observed as stocks with different sizes in the blue-augen gneisses, at the vicinity of Birgi town. The mineral composition of these unmetamorphosed basic plutons consists of "Plagioclase + orthopyroxene + clinopyroxene + olivine + biotite + apatite + zoisite + zircon and +garnet". The extrusive rocks which are the volcanic equivalents of these basic and acidic plutons are observed at two different parts of the area. According to their mineralogical and chemical composition, these volcanic rocks are of andesite and basalt in composition. All the rock units exposed at the investigated area are unconformably covered by the Neogene aged unmetamorphosed sedimentary rocks.

### PETROGRAPHY OF THE LEPTITES

Leptites show a widespread distribution in the Ödemiş submassif, the middle part of the Menderes massif. These rocks which occur between the basal gneiss and schist vary in thickness from 20-25 m to several kilometers. Purple, grayish and gray coloured leptites are hard, massive and generally badly schistosed rocks. These metavolcanites are often cut by the pegmatitic veins, ranging from mm to m in thickness. These pegmatitic veins are composed of feldspar, quartz, muscovite and tourmaline. In the field, one of the characteristic features of the leptites is the presence of the old sills and dykes which exhibit hornfelsic texture and rich-in pyroxene and anorthite (Kun and Candan, 1987). These gabbroic vein rocks were simultaneously metamorphosed with the leptites.

The locally preserved old porphyritic texture can be observed in the leptites. This evidence clearly suggests that the origin of these rocks called as leptites in this paper, is volcanic. In these porphyritic rocks which show a foliated andesitic appearance, the old phenocrysts were oriented by the metamorphism (Fig. 3). These phenocrystal relicts were filled by some light coloured minerals such as quartz, muscovite, feldspar, sillimanite, and gamet.



**Fig. 3 - Photograph of the relict porphyritic texture of the primary volcanics observed in leptites.**

The field observations show that leptites occur as a guide horizon between the basal gneiss complex which were derived from graywackes, and schists in the Menderes massifs. But, because of the presence of the blue-augen gneisses most probably of volcanic origin like leptites the contact mapped between the basal gneiss complex and leptite, does not correspond to the primary graywacke/volcanic rock boundary (Fig. 4).




Fig. 4 - Cross-section showing the contact relation between blue augen gneiss and leptite.

From the petrographic studies, it has been observed that the leptites exhibit a typical fine-grained, nondirectional granoblastic and hornfelsic texture (Fig. 5). But some samples, especially those which are rich in biotite, may show a poorly developed schistosity. The general mineral compositions of the leptites consist of "Quartz-plagioclase-orthoclase-garnet-biotite-muscovite-epidote-sillimanite-kyanite-tourmaline-zircon and apatite". The leptites can be divided into three major groups according to their macro and micro features: 1 - Garnet-biotite leptites; 2 - Sillimanite-kyanite leptites; 3 - Spotted-leptites.




Fig. 5 - Photomicrograph showing the typical fine-grained, nondirectional hornfelsic texture in leptites. 2.5 X, crossed nicols. Q: Quartz, Plj: Plagioclase.

#### 1 - Garnet-biotite leptites

The great majority of the leptites exposed in the study area belongs to this group. While the percentage of garnet reaches up to 35 % and biotite up to 40 %, muscovite is always present in the amounts of less than 5 % in these rocks. Garnet-biotite leptites which exhibit a poorly developed schistosity consist mainly of quartz, orthoclase, plagioclase, biotite, muscovite, garnet, epidote, zoisite, apatite, tourmaline and zircon.




Fig. 6 - Sillimanite fibrolites occurring at the contacts of the feldspars. Sil: Sillimanite, 10 X, plane light.

## 2 - Sillimanite-kyanite leptites

This type of leptites closely resemble the garnet-biotite leptites. The distribution of the sillimanite-kyanite leptites in the investigated area may be only identified by the microscopic studies. With the increasing percentage of the sillimanite content, these rocks exhibit a harder and rougher appearance. The percentage of the sillimanite ranges from 1 % to 15 % (Fig. 6). But, the kyanite content of these leptites does not exceed 3 %. The formation of the sillimanites can be divided into two groups by their textural evidences. 1) Sillimanites, which occur as fibrolites at the contacts of the feldspars, and 2) Sillimanites, which replace the garnet and biotite crystals (Fig. 7).



Fig. 7 - Sillimanite crystals occurring by the replacement of the biotites. Sil: Sillimanite; Bio: Biotite; 10 X, plane light.



### 3 - Spotted leptites

These leptites, which were rarely observed can be easily recognized in the field by their white spots. These appearance are caused by the preserved relict porphyritic textures of the primary volcanic rocks, and are of great importance to clarify the parent rocks of the leptites. These spots were generally filled by the light coloured minerals such as quartz, feldspar and muscovite. In general, an euhedral garnet crystal is situated at the center of these voids and this core is rimmed by the sillimanite and biotite zones (Fig. 8). The general mineral compositions of the spotted leptites show a great similarity to the sillimanite-kyanite leptites and consist of "Quartz-orthoclase-plagioclase-muscovite-biotite-sillimanite-kyanite-gamet-zircon and apatite".




Fig. 8 - Photomicrograph of the white-spots in the leptites. An euhedral garnet crystal is situated at the center of these voids. Sil: Sillimanite; Gr: Garnet; 10 X, plane light.

### CHEMISTRY OF THE LEPTITES

In general, the leptites exhibit more or less a homogeneous mineral composition. But, of course, the amount of garnet, biotite and sillimanite in the rocks affect the chemical composition of the leptites. The average contents of SiO<sub>2</sub>, Al<sub>2</sub>O<sub>3</sub>, FeO, CaO, MgO, Na<sub>2</sub>O and K<sub>2</sub>O in the leptites can be given as 61-74 %, 11-18 %, 4-6 %, 0.7-2.6 %, 1-3 %, 1.64.3 % and 2-4.5 % respectively. TiO<sub>2</sub> is always present in amounts of less than 1 % (Table 1). A comparison of the Sweden (Löfgren, 1979) leptites with those of the study area in terms of SiO<sub>2</sub> contents is given in Table 2. The leptites from Sweden are divided into 6 different groups. But, the leptites in Ödemiş submassif of the Menderes massif show only 3 groups and don't contain the extreme acidic and basic groups.

The chemical analysis were plotted on several diagrams to find out the properties of the original volcanics of the leptites. In order to eliminate the influences of possible element migrations during the high grade metamorphism, in addition to the major oxides, the diagrams based on the trace elements were also used.

In the Na<sub>2</sub>O+K<sub>2</sub>O/SiO<sub>2</sub> diagram proposed by Zanettin-(1984), the leptites are concentrated in rhyolite, dacite and andesite areas (Fig. 9). In Kistler and Evernden (1971), Winchester and Floyd (1977) diagrams based on the Rb vs Sr and SiO<sub>2</sub> % versus Zr/TiO<sub>2</sub> respectively, all the examined metavolcanics plot in the fields of andesite, rhyodacite and rhyolite (Fig. 10 and 11).

**Table 1 - Comparison of the Sweden and Ödemiş submassif leptites in terms of SiO<sub>2</sub> contents**

	Group 1 % SiO <sub>2</sub> 50-55	Group 2 % SiO <sub>2</sub> 55-60	Group 3 % SiO <sub>2</sub> 60-65	Group 4 % SiO <sub>2</sub> 65-70	Group 5 % SiO <sub>2</sub> 70-75	Group 6 % SiO <sub>2</sub> 75-80
<b>Leptites of Sweden</b>						
SiO <sub>2</sub>	53.66	57.45	62.82	67.20	72.64	77.28
Al <sub>2</sub> O <sub>3</sub>	18.74	16.49	15.92	14.67	13.06	12.00
TiO <sub>2</sub>	.93	1.06	.59	.52	.26	16
Fe <sub>tot</sub>	8.89	8.33	5.08	4.34	2.60	1.35
MgO	4.18	3.54	2.58	1.81	1.14	80
CaO	5.70	5.06	3.27	2.10	1.36	70
Na <sub>2</sub> O	2.86	2.58	2.58	2.46	2.42	3.93
K <sub>2</sub> O	3.36	2.93	3.96	4.94	4.90	2.97
<b>Leptites of the Ödemiş submassif</b>						
SiO <sub>2</sub>			63.38	65.70	71.68	
Al <sub>2</sub> O <sub>3</sub>			15.27	15.01	13.09	
TiO <sub>2</sub>			.88	.80	.73	
Fe <sub>tot</sub>			5.40	4.77	3.92	
MgO			2.58	1.98	1.67	
CaO			1.55	1.54	1.33	
Na <sub>2</sub> O			3.19	3.02	3.24	
K <sub>2</sub> O			3.34	3.11	2.62	




Fig. 9- Na<sub>2</sub>O + K<sub>2</sub>O versus SiO<sub>2</sub> diagram after Zanettin (1984).




Fig. 10 - Rb (ppm) - Sr (ppm) variation diagram after Kistler and Evernden (1971).




Fig. 11 - SiO<sub>2</sub> (Wt %) vs Zr (ppm) / TiO<sub>2</sub> (Wt %) diagram after Winchester and Floyd (1977).

In order to determine the kindred of the primary volcanics, several diagrams based on major oxides are used. In the Rittmann (1962) diagram, all the leptites are accumulated in the field of calc-alkaline series (Fig. 12). In another diagram using the same oxides, the great majority of the rocks are concentrated in the calc-alkaline series but only a few samples extended into the tholeiitic area (Fig. 13). In the Miyashiro (1975) diagram, where FeO\*/MgO versus SiO<sub>2</sub> is plotted the samples fall again in the calc-alkaline field (Fig. 14).

	O-305	§-206/A	§-51	§-178	O-202/A	O-118	O-226	O-316/A	O-292	O-301/C
SiO <sub>2</sub>	69.85	71.28	69.24	65.69	70.10	64.01	72.08	71.69	71.57	68.60
Al <sub>2</sub> O <sub>3</sub>	14.23	13.36	14.60	16.26	13.74	16.88	13.16	13.96	14.14	15.00
TiO <sub>2</sub>	.69	.65	.65	.76	.81	.78	.63	.60	.54	.68
Fe <sub>2</sub> O <sub>3</sub>	1.59	2.02	1.87	4.52	2.25	3.37	.99	3.90	1.54	2.79
FeO	2.78	1.96	2.54	1.52	2.17	2.54	2.70	.00	1.90	1.96
MnO	.05	.06	.07	.07	.05	.08	.06	.04	.06	.05
MgO	1.86	1.43	1.78	2.61	1.58	2.54	1.46	1.36	1.35	1.91
CaO	1.09	1.44	1.57	.70	.95	.97	1.16	.76	.98	.83
Na <sub>2</sub> O	2.32	2.79	3.63	1.69	2.74	2.33	3.35	2.40	4.07	3.02
K <sub>2</sub> O	3.18	3.36	3.08	4.07	3.08	3.84	2.31	2.60	2.20	3.11
P <sub>2</sub> O <sub>5</sub>	14	18	18	.16	.17	.20	.12	.12	.18	.17
IL <sub>2</sub> O	1.36	1.36	.93	.95	1.37	1.92	.85	2.02	.84	1.53
<b>Total</b>	<b>99.14</b>	<b>99.89</b>	<b>100.14</b>	<b>99.00</b>	<b>99.03</b>	<b>99.46</b>	<b>98.96</b>	<b>99.45</b>	<b>99.37</b>	<b>99.65</b>
Zn	73	50	0	128	60	125	51	52	48	70
Ni	61	54	56	40	47	66	45	57	43	49
Cu	3	3	5	24	7	13	6	12	15	14
Cr	120	124	114	133	152	144	125	120	104	125
V	93	86	89	106	100	115	89	82	67	90
Zr	193	238	183	186	337	171	224	206	183	205
Y	23	22	23	24	25	27	21	23	19	21
Sr	192	196	222	131	209	155	258	208	231	201
Rb	118	88	97	133	111	124	78	89	107	102
Hf	8	8	6	8	9	6	10	6	7	8
	A-39	A-18A	§-209/B	G-22	O-203/A	O-268	G-40	480/B		
SiO <sub>2</sub>	70.37	71.58	69.12	66.90	70.40	69.25	64.82	67.73		
Al <sub>2</sub> O <sub>3</sub>	15.06	13.65	14.64	16.01	13.83	14.14	16.63	14.40		
TiO <sub>2</sub>	.67	.78	1.01	.72	.84	.70	.84	.70		
Fe <sub>2</sub> O <sub>3</sub>	3.64	5.03	5.16	6.37	4.78	5.30	6.75	5.18		
MgO	.96	1.43	1.50	1.76	1.43	1.54	1.91	2.75		
CaO	1.87	1.72	2.23	1.04	2.61	2.26	.89	1.64		
Na <sub>2</sub> O	3.30	3.75	3.53	2.12	4.00	3.59	2.21	4.56		
K <sub>2</sub> O	4.33	2.33	2.51	3.76	2.17	2.82	4.16	2.33		
K.K.	.24	.37	.28	1.19	.22	.38	1.90	.72		
<b>Total</b>	<b>100.44</b>	<b>100.64</b>	<b>99.98</b>	<b>99.87</b>	<b>100.28</b>	<b>99.98</b>	<b>100.11</b>	<b>99.96</b>		
Ba	653.3	305.0	921.9	637.0	468.8	510.4	694.9	281.5		
Nb	8.9	10.5	15.5	19.2	14.9	11.6	13.6	13.5		
Zr	128.0	191.6	122.6	154.9	139.1	125.6	130.4	156.9		
Y	19.7	13.7	27.7	14.0	2.5	E	27.2	11.8		
Sr	115.5	226.2	480.0	145.7	305.6	260.2	164.3	38.0		
Rb	128.8	79.5	64.0	124.3	68.1	96.5	126.2	65.1		
	448	431	486	463/A	§-24	§-22/D	463/A	§-101	§-44	§-63
SiO <sub>2</sub>	71.66	74.23	71.94	70.99	65.87	71.06	63.30	71.33	72.67	69.02
Al <sub>2</sub> O <sub>3</sub>	12.24	11.71	12.91	13.18	16.76	13.93	16.94	14.12	13.28	14.50
TiO <sub>2</sub>	.66	.73	.75	.76	.88	.70	.86	.64	.71	.85
Fe <sub>2</sub> O <sub>3</sub>	4.21	4.36	4.39	4.56	5.08	4.21	5.97	4.03	3.77	4.21
MgO	2.28	1.65	1.72	1.64	1.86	1.43	2.08	1.43	1.34	1.47
CaO	.99	.82	1.96	1.21	.70	.84	2.66	1.16	.97	2.27
Na <sub>2</sub> O	3.47	2.76	3.34	2.90	2.54	2.91	4.22	3.57	3.17	4.17
K <sub>2</sub> O	3.19	2.27	2.24	2.60	3.85	2.99	2.43	2.58	2.74	2.27
K.K.	.85	1.22	.54	1.27	1.94	1.64	1.50	1.02	1.12	1.02
<b>Total</b>	<b>99.55</b>	<b>99.75</b>	<b>99.79</b>	<b>99.11</b>	<b>99.48</b>	<b>99.71</b>	<b>99.96</b>	<b>99.88</b>	<b>99.77</b>	<b>99.78</b>
Ba	563.9	423.2	451.6	598.3	618.3	707.6	468.2	594.2	609.6	485.6
Nb	10.8	13.6	15.2	5.3	18.9	—	12.9	13.4	11.7	8.8
Zr	111.8	148.4	141.7	126.5	151.5	—	137.4	149.4	180.8	266.9
Y	14.1	20.7	6.6	19.4	27.7	—	13.9	21.4	17.1	14.0
Sr	179.0	191.9	182.0	180.2	181.3	—	279.2	247.7	246.5	34.9
Rb	81.6	77.4	73.0	79.7	104.1	—	94.5	80.7	81.2	77.2
	465/C	467	449	432/A	463/D	426	437/B	380	469	363
SiO <sub>2</sub>	73.16	64.82	72.03	71.51	71.50	69.65	63.04	71.29	72.76	66.56
Al <sub>2</sub> O <sub>3</sub>	10.68	15.15	12.99	12.96	12.69	12.99	16.19	12.19	11.97	14.27
TiO <sub>2</sub>	.98	.81	.59	.70	.72	.90	1.09	.84	.79	1.15
Fe <sub>2</sub> O <sub>3</sub>	5.25	5.37	3.94	3.95	4.54	4.47	6.04	4.57	4.94	6.02
MgO	2.29	3.07	2.11	2.14	2.35	2.32	3.17	2.40	1.41	2.17
CaO	1.68	2.01	1.71	1.28	1.00	3.16	2.81	1.54	1.34	2.35
Na <sub>2</sub> O	2.99	4.31	3.52	4.22	3.11	3.55	3.99	3.50	2.94	3.08
K <sub>2</sub> O	2.18	2.43	2.35	2.16	2.58	2.31	2.46	2.35	2.33	2.77
K.K.	.59	1.53	.85	.76	1.27	.85	.80	1.05	1.32	1.22
<b>Total</b>	<b>99.80</b>	<b>99.58</b>	<b>100.09</b>	<b>99.68</b>	<b>99.79</b>	<b>100.20</b>	<b>99.59</b>	<b>99.73</b>	<b>99.80</b>	<b>99.59</b>
Ba	655.6	363.6	426.3	413.3	820.0	442.3	639.4	530.7	528.2	604.0
Nb	12.3	13.6	12.9	14.9	11.6	13.6	—	10.1	12.8	5.8
Zr	282.1	116.3	117.1	104.9	131.3	161.5	—	190.6	214.1	188.0
Y	17.0	16.7	12.1	14.8	10.0	21.0	—	114.4	17.9	14.4
Sr	172.1	143.0	182.3	203.2	183.3	405.7	—	35.1	37.9	257.4
Rb	57.3	109.1	79.1	78.5	76.5	52.9	—	76.4	75.2	94.9
	482/B	391	§-59	432/B	447	O-316	463/B	§-102	§-167	467
SiO <sub>2</sub>	66.61	71.79	67.27	69.30	68.06	65.40	61.62	62.96	66.94	64.15
Al <sub>2</sub> O <sub>3</sub>	15.74	12.14	16.05	13.82	14.00	17.04	18.26	16.94	15.12	17.58
TiO <sub>2</sub>	.82	.86	.82	.81	.77	.77	.95	.91	.85	.80
Fe <sub>2</sub> O <sub>3</sub>	6.26	5.32	5.18	4.75	5.53	5.53	5.97	6.41	5.09	6.65
MgO	2.16	1.63	1.74	1.76	1.86	1.79	2.30	2.47	1.54	2.95
CaO	1.21	1.18	.80	2.58	1.75	.80	1.88	.65	2.13	.29
Na <sub>2</sub> O	2.94	2.39	2.60	3.14	3.44	2.48	4.48	2.62	3.11	2.85
K <sub>2</sub> O	3.30	2.71	3.86	2.86	2.75	3.42	3.07	4.50	3.24	3.16
MnO	.06	.07	.04	.05	.05	.05	—	—	.07	.04
K.K.	1.13	1.26	1.49	.80	1.23	2.69	.83	.39	1.29	1.25
<b>Total</b>	<b>100.23</b>	<b>99.37</b>	<b>99.85</b>	<b>100.07</b>	<b>99.44</b>	<b>99.97</b>	<b>99.36</b>	<b>99.85</b>	<b>99.38</b>	<b>99.72</b>
Ba	952.8	583.4	624.9	715.9	536.0	599.0	857.2	710.6	539.09	—
Nb	11.8	13.7	12.0	—	10.7	14.1	14.3	15.6	11.6	—
Zr	76.6	240.0	133.0	—	103.2	122.7	123.9	125.1	172.0	—
Y	10.6	17.0	22.6	—	12.0	13.8	33.5	40.3	25.3	—
Sr	157.7	172.5	169.9	—	266.7	189.2	275.4	182.3	134.2	—
Rb	109.9	85.1	118.0	—	128.3	114.5	101.9	142.7	102.7	—




Fig. 12 - Place of the Ödemiş leptites on Rittmann (1962) diagram.




Fig. 13 - Place of the Ödemiş submassif leptites on alkali-silica diagram.




Fig. 14 - FeO\* / MgO versus SiO<sub>2</sub> diagram after Miyashiro (1975).

In the log  $\tau$  versus log  $\delta$  diagram suggested by Gottini (1968), all the samples fall in the sialic origin (Fig. 15). There are several diagrams to explain the tectonic environment of the volcanics on the light of the plate tectonic theory. In the Morrison (1980) diagram, where K<sub>2</sub>O are plotted against SiO<sub>2</sub>, all the samples lie within the calc-alkaline island arc fields (Fig. 16). In the TiO<sub>2</sub> versus FeO/MgO diagram advanced by Glassley (1974), the leptites of Ödemiş submassif are concentrated in the island arc field too (Fig. 17).




Fig. 15 - Log  $\tau$  vs log  $\delta$  diagram after Gottini (1969).




Fig. 16 - Variation diagram of K<sub>2</sub>O contents with SiO<sub>2</sub> after Morrison (1980).




Fig. 17 - TiO<sub>2</sub> versus FeO\* / MgO diagram after Glassley (1974).

The  $Ti \times 10^{-2} / Zr / Y \times 3$ ,  $Ti \times 10^{-2} / Zr / Sr \times 2$  and  $Ti / Zr$  diagrams proposed by Pearce and Cann (1973) were applied to the leptites and it has been seen that the original rocks of the leptites were the calc-alkaline island arc volcanics (Fig. 18). The analysis of the leptites were plotted on the  $TiO_2$  (%) versus Zr (ppm) diagram which was used by Gass (1982) to find out the tectonic environment of the metavolcanics belonging to the Upper Pan-African orogenesis (Fig. 19). It may be concluded from this diagram that the Menderes massif leptites show great similarity with the metavolcanics exposed at the NE Africa and SW Arabian Peninsula.

Finally, it can be suggested that the parent rocks of the leptites in the Ödemiş submassif of the Menderes massif are dacitic and rhyolitic volcanics. These island arc volcanics have sialic origin and dominant calc-alkaline kindred.




Fig. 18 -  $Ti \times 10^{-2}/Zr/Y \times 3$ ,  $Ti/100/Zr/Sr/2$  and  $Ti/Zr$  diagrams after Pearce and Cann (1973). A = A and B: Low K-Tholeiites, C and B: Calc-alkaline basalts, B: Ocean floor basalts, D: Within plate basalts. B = A: K-Poor tholeiites, B: Ocean floor basalts, C: Calc-alkaline basalts, C = D and B: Ocean floor basalts; A and B: Low K-Tholeiites, C and B: Calc-alkaline basalts.




Fig. 19 - Place of the leptites of Ödemiş submassif on the  $TiO_2$  (%) /  $Zr$  (ppm) diagram.



## DISCUSSION AND CONCLUSIONS

The general evolution of the Menderes massif and the position of the leptites (metavolcanics) which are situated at the Ödemiş submassif can be briefly summarized as follows:

According to the recent studies in the Menderes massif, the initial sedimentation age of the gneisses occurring at the lower level of the metamorphic sequence is about 545-670 Ma old (Satir ve Friedrichsen, 1986). These sediments which were widely composed of graywackes were subjected to a high grade metamorphism about 500 Ma ago and were cut by the tonalitic-granitic intrusions about 470 Ma ago (Satir and Friedrichsen, 1986). In most places, these gneissic basement was covered by the primary volcanic rocks of the leptites which were the surface equivalents of these post-metamorphic acidic plutons (Kun and Candan, 1987; Dora et al., 1988). These volcanics, rhyolite and dacite in-composition, are the initial rocks of the leptites. In Ordovician, the sedimentation was started in the region and until the Paleocene, sedimentary series which was composed of the flysh type sediments and platform type limestones were deposited on these volcanics (Dürr 1975; Çağlayan et al., 1980; Konak et al., 1987; Dora et al., 1989). The last main metamorphism which was affected the Menderes massif has occurred in Upper Paleocene-Lower Eocene time (Dürr, 1975; Dürr et al., 1978; Şengör et al., 1984; Kun and Candan, 1987; Andreissen et al., 1979). The leptites in Ödemiş submassif which are of acidic in-composition were formed by this high grade metamorphism (Kun and Candan, 1987).

In recent years, it has been claimed that the Menderes massif is part of the Pan-African continental crust exposed in NE Africa and Arabian Peninsula (Şengör et al., 1984; Dora et al., 1988). According to Gass (1982), the Upper Pan-African orogenesis in these regions is especially characterized by the silicic magmatic activity and acidic volcanics which vary in composition from andesite to rhyolite. These island arc type volcanic rocks have a calc-alkaline kindred.

It follows that the leptites (metavolcanics) exposed in the Ödemiş submassif of the Menderes massif shows a great similarity with the Upper Pan-African volcanics in NE Africa in terms of chemical composition and geotectonic position. It may be suggested by these chemical and geological evidences that the leptites of the Ödemiş submassif represent the North-western extension of the island arc volcanic belt widely occurring in NE Africa and Arabian Peninsula, in relation to the late phases of the Pan-African orogenesis.

## ACKNOWLEDGEMENTS

This work was supported by TÜBİTAK (TBAG-688). We also thank Mualla Gürle for considerable drafting assistance.

*Manuscript received February 26, 1990*

## REFERENCES

- Andriessen, P.A.; Boerlijck, N.A.I.M.; Hebeda, E.H.; Priem, H.N.A.; Verdunman, E.A. Th. and Verschure, R.H., 1979, Dating the events of metamorphism and granitic magmatism in the Alpine Orogen of Naxos (Cyclades, Greece): *Contrib. Mineral. Petrol.*, 69, 215-225.
- Çağlayan, M.A.; Öztürk, E.M.; Öztürk, Z.; Sav, H. and Akat, U., 1980, Menderes Masifi güney kanadına ait bulgular ve yapısal yorum: *Jeo. Müh. Der.*, 9-17.
- Dora, O. Ö.; Kun, N. and Candan, O., 1988, Metavolcanics (leptite) in the Menderes Massif: A possible paleoarc volcanism: *ODTÜ Melih Tokay Sempozyumu Bull.*, (in print).
- ; — and —, 1989, Menderes Masifi'nin metamorfik tarihçesi ve jeotektonik konumu: Ç.Ü. Ahmet Acar Jeoloji Sempozyumu, *Bildiri Özetleri Kitapçığı*, p. 5.
- Dürr S., 1975, Über alter und geotektonisch stellung des Menderes Kristallins/SW - Anatolien und seine sequivalente in dermittleren Aegaeis: *Marburg/Lahn*, 107 p.

- Dürr, S.; Alther, R.; Keller, J.; Okrusch, M. and Seidel, E., 1978, Alps, Apennines, Hellenides: Inter-Union Commission on Geodynamics Scientific Report No: 38, p. 454-477.
- Gass, G., 1982, Upper Proterozoic (Pan African) calc-alkaline magmatism in north-eastern African and Arabia: In andesite, cd. Thorpe, R.S., John Willay and Sons.
- Glassley, W., 1974, Geochemistry and tectonics of the Crescent volcanic rocks, Olympic peninsula: Geol. Soc. Amer. Bull., 85, 785-794, Washington.
- Goltini, V., 1968, The TiO<sub>2</sub> frequency in volcanic rocks: Geol. Rdsch., 168 pp.
- Kistler, R.W. and Evenden, J.F., 1971, Sierra Nevada Plutonics cycle: Part I, origin of composite granitic batholiths: Geol. Soc. Amer. Bull., 82, 853-868.
- Konak, N.; Akdeniz, N. and Öztürk, E.M., 1987, Geology of the south of Menderes Massif: IGCP Project No. 5, 42-53, Turkey.
- Kun, N., 1983, Cine dolayının petrografisi ve Menderes Masifinin güney kesimine ait petrolojik bulgular: Doktora tezi, DEÜ, İzmir.
- and Candan, O., 1987, Ödemiş Asmasifindeki leptitlerin dağılımı, konumları ve oluşum koşulları: TBAG-688 number of project, 133 P.
- and ———, 1988, Menderes Masifindeki Erken Paleozoik yaşlı bazik damar kayaları: Hacettepe Üniversitesi Yerbilimleri Derg. (in print).
- Löfgren, C., 1979, Do leptites represent Precambrian island arc rocks: Lithos, 12, 159-165.
- Miyashiro, A., 1975, Volcanic rock series and tectonic setting: In annual review of earth and planetary Science, 3, 51-269.
- Morrison, G. W., 1980, Characteristics and tectonic setting of the shoshonite rocks association: Lithos, 13, 97-108.
- Pearce, J.A. and Cann, J.R., 1973, Tectonic setting of basic volcanic rocks determined using trace element analysis: Earth planet, Sci. Lett 19, 290-300.
- Phillipson, A., 1911, Reisen und forschungen im Westlichen Kleinasien: Petermanns Mitt Erganzonpsheft, 172, Gotha.
- Rittmann, A., 1962, Volcanoues and their activity: John Wiley and Sons, Newyork, London, 305 pp.
- Satr, M. and Friedrichsen, H., 1986, The origin and evolution of the Menderes Massif, W-Turkey: A rubidium/strontium and oxygen isotope study, Geol. Rundschau 75/3, 703-714.
- Schuilng, R.D., 1962, Türkiye'nin güneybatısındaki Menderes migmatit kompleksinin petrolojisi, yaşı ve yapısı hakkında: MTA Bull. 58, 71-84, Ankara.
- Şengör, M.C.; Saur, M. and Akkök, R., 1984, Timing of tectonic events in the Menderes Massif, Western Turkey: Implication for tectonic evolution and evidence for Pan-African basement in Turkey: Tectonics, Vol. 3, no. 7, 693-707.
- Winchester, J.A. and Floyd, P.A., 1977, Geochemical discrimination products using immobile elements: Chem. Geol. 20, 325-343.
- Zanettin, B., 1984, Proposed new chemical classification of volcanic rocks: Episodes, 714, 19-20.